



TITLE:

Chemical, Mineralogical and Micromorphological Properties of Glaebules in Some Tropical Lowland Soils

AUTHOR(S):

Furukawa, Hisao; Handawella, James; Kyuma, Kazutake; Kawaguchi, Keizaburo

CITATION:

Furukawa, Hisao ...[et al]. Chemical, Mineralogical and Micromorphological Properties of Glaebules in Some Tropical Lowland Soils. 東南アジア研究 1976, 14(3): 365-388

ISSUE DATE:

1976-12

URL:

<http://hdl.handle.net/2433/55851>

RIGHT:

Chemical, Mineralogical and Micromorphological Properties of Glaebules in Some Tropical Lowland Soils

Hisao FURUKAWA*, James HANDAWELLA**,
Kazutake KYUMA***, and Keizaburo KAWAGUCHI*

Introduction

In the course of field studies on padi soils in South and Southeast Asia, different kinds of secondary formations embedded in and/or resting on the soils were found under a wide variety of geomorphological and geological settings. These secondary formations correspond to what Kellogg (1949) called "concretions or nodules in a matrix of unconsolidated material", and "consolidated mass of such concretions or nodules". They are comprehensively called glaebules in this paper according to Brewer's nomenclature (1964).

Some properties of these glaebules from various soil samples were studied to obtain a general view on the prevailing conditions in soils and sediments which give rise to glaebules.

The geomorphological setting relative to ferruginous glaebule formation in the Central Plain of Thailand was previously described by Takaya (1968). He found the most striking occurrence of pisolitic concretions in the soil cover on Terrace II (Upper Pleistocene) where the soils are mostly of low humic gley nature (Moormann and Rojanasoonthon, 1967). Terrace III and the Peneplain were described as capped by thin laterite and thick hard laterite respectively. Takaya presumed these glaebules to be formed *in situ*. The length of time of weathering to which each geomorphological surface was exposed was thought to be relevant to the occurrence of the various types of glaebules.

Fridland (1964) reported that pisolitic concretions occur on the middle terraces but not on the low terraces and flood plains of the Ngan-Fo river of Vietnam.

Although these findings give useful field criteria for classifying geomorphological surface and for correlating specific stratigraphic strata in the quaternary deposits of different regions, the occurrence of a certain type of glaebule can not necessarily be used to estimate the age of soil material, even if it may be used to estimate the areal

* 古川久雄, 川口桂三郎, Department of Soil Science, Faculty of Agriculture, Kyoto University.

** Agriculture Research Station, Maha, Ilupallama, Sri Lanka.

*** 久馬一剛, The Center for Southeast Asian Studies, Kyoto University.

extent of a soil with the same morphology. The two differently evolving types of terrain distinguished by geomorphologists should be borne in mind. Van Wambeke (1962) has clearly illustrated the distribution pattern of soil units in the two different types of geomorphological process; one of these processes is characterized by rapidly rejuvenating erosion cycles, resulting in terrace formation, and the other by the development of a single erosion surface. Old soils are found on elevated terraces in the former, and on lower-lying sites in the latter.

Terrains in tropical Asia are considered to have been affected by both processes; soils on upper terraces are in general more weathered and mature than lower-lying soils, but at the same time they are more likely to have received soil material from rapidly eroding hillslopes.

Accordingly, the glaeboles most frequently found in soils on older terraces may be partly relicts, and partly present formations resulting from the modern pedo- and geo-genetic environments. These considerations necessitate a detailed description of the properties of the glaeboles, taking care not to make hasty conclusions on the pedological and geomorphological significance of the glaeboles. In this paper we intend to relate the pedogenetic conditions in the field to the properties of the glaeboles.

Materials and methods

All the glaebole samples from tropical Asia come from padi soils, with a few manganiferous glaeboles from Reddish-brown soil on coral limestone (Matsuzaka *et al.*, 1971) from Okinawa Prefecture, Japan, and from Terra roxa estruturada eutrofica on basic eruptiva from Brazil. An indurated vesicular laterite crust derived from saprolitic agrillaceous rocks in West Malaysia was also included in the samples.

The distribution of sample sites is mapped in Fig. 1 in relation to Kyuma's climatic regional division (1972).

Indurated thick hard laterite is widely distributed in regions II, IV, V, and VII, and pisolitic ferruginous glaeboles seem to occur predominantly in regions V and VII where the climate is characterized by the presence of dry season, namely the tropical monsoon (region V) and tropical-subtropical monsoon (region VII) climatic regions.

Soils rich in bases and montmorillonitic clay, such as those occurring in central Luzon in the Philippines, contain glaeboles of manganiferous nature. Ferruginous glaeboles are also found on older alluvium (Okagawa, 1970), but their size, roundness, and greasy surface texture are not as developed as that of glaeboles occurring in the central plain of Thailand and in India.




Fig. 1 Glaebule sampling sites (using Kyuma's climatic regional division, 1972)

The infrequent occurrence of ferruginous glaebules in the soils of Java is also noteworthy, and may be related to the continuous rejuvenation of soil parent materials through active volcanism.

The distribution of carbonate glaebules is related to calcium-rich geology, such as leucite-bearing eruptiva and limestone.

The distribution of glaebules in the soil profile is variable and does not appear to show any regular change in abundance, size, and hardness related to position within a profile. A certain regularity, was noted in some previous studies, reviewed by Drosdoff and Nikiforoff (1940), in soils in the temperate regions, and, in a study by Nye (1955), in West African soils. All these soils can be regarded as more or less sedentary, while the samples used in the present study are transported soils.

Soil units at the sampling sites were given approximated names, using several sources (Moormann and Rojanassonthon, 1967; Survey of India Offices, 1954; Suprptoahardjo *et al*, 1960; Takaya and Kyuma, 1968; Barrera, 1964).

Glaebules samples were ground by a spexmill or an agate mortar. Major chemical composition of the samples was determined by means of the X-ray fluorescence spectroscopy, in which ignited ground powder was fused into a glass disc with a flux ($\text{Li}_2\text{B}_4\text{O}_7$ - Li_2CO_3 - La_2O_3 mixture) according to the method described by Norrish and Chappell (1967).

The calibration curves drawn with synthetic standard mixtures were sufficiently

straight for a wide range of concentrations. The mean total of nine oxides (SiO_2 , Al_2O_3 , Fe_2O_3 , Mn_3O_4 , CaO , MgO , K_2O , and P_2O_5) was 97.0 ± 5.0 (57 samples) on an ignited basis. The content of each oxide was recalculated to total 100%, neglecting the matrix correction which would have yielded a small change in content.

For mineralogical composition the ground powder was mounted on etched glass slides and scanned by an X-ray diffractometer with Fe K_α . Quartz, feldspar, goethite, haematite, calcite, and sometimes clay minerals were recognized and estimated semi-quantitatively by the peak height.

Oriented clay specimen obtained by Mehra-Jackson's deferration treatment (1960) was X-ray diffracted, and the clay mineral composition of the glaeble samples was compared with that of the soil matrix.

For micromorphological observation, glaebles embedded in polyester resin under vacuum were cut and ground to thin sections by conventional methods in geological laboratories. The thin sections were studied under transmitted and reflected light using a polarizing microscope. Color and fabric of the groundmass, voids, and skeletal grains were observed, described and interpreted mainly using the terms proposed by Kubiena (1970).

Results and discussions

1. Appearance and chemical composition

The glaebles studied may be grouped into three according to chemical composition, that is, ferruginous, manganiferous, and carbonate glaebles (Table 1).

The appearance of the glaebles, described in terms of color, size, shape, surface texture, surface lustre, hardness, etc. is widely variable (Fig. 2). Ferruginous glaebles, in particular, showed wide variation;

- (1) dark brown to brown, spherical granular, nodular to botryoidal of variable size, with smooth surface texture and greasy lustre,

Table 1 Mean chemical composition of various glaebles (% on ignited basis)

	SiO_2	Al_2O_3	Fe_2O_3	Mn_3O_4	CaO	MgO	TiO_2	K_2O	P_2O_5	$\text{SiO}_2/\text{Al}_2\text{O}_3$
Ferruginous glaebles with low $\text{SiO}_2/\text{Al}_2\text{O}_3$ (subspherical-blocky)	20.0	18.5	58.5	0.33	0.08	0.73	0.82	0.18	0.37	1.85 ± 0.13 ($n=13$)
Ferruginous glaebles with high $\text{SiO}_2/\text{Al}_2\text{O}_3$ (spherical)	27.8	10.7	57.4	1.65	0.49	0.80	0.64	0.27	0.21	4.60 ± 1.43 ($n=21$)
Manganiferous glaebles with smooth surface (spherical)	36.9	11.5	32.2	20.8	0.72	0.87	1.39	0.16	0.10	4.90 ± 1.42 ($n=6$)
Manganiferous glaebles with mammilated surface (oblate)	28.1	17.4	23.0	26.1	1.26	1.07	1.75	0.87	0.39	2.80 ± 0.39 ($n=5$)
Carbonate glaebles (warty-nodular)	29.7	7.1	9.3	3.68	45.6	3.01	0.41	0.89	0.19	7.40 ± 1.60 ($n=8$)




Fig. 2 Representative glaebules and indurated laterite

1. Manganiferous glaebules from Reddish-brown soil on coral limestone, Okinawa Prefecture, Japan. Slightly mammilated surface texture.
2. Manganiferous glaebules from a Latosol on intermediate to basic eruptiva, Luzon, the Philippines. Smooth surface texture.
3. Warty blocky carbonate glaebules, coexisting with spherical ferruginous glaebules with greasy lustre, and subspherical manganiferous glaebules. Sampled on old alluvium in Central Thailand.
4. Botryoidal ferruginous glaebules forming a hardpan. Sampled on old alluvium in Northeast Thailand.
5. Spherical ferruginous glaebules with smooth surface texture, but with dull lustre. Sampled on old alluvium in intermountain basin of Northern Thailand.
6. Spherical ferruginous glaebules with bleached appearance. Sampled from a Low Humic Gley soil in Central Thailand.
7. Nodular and spherical ferruginous glaebules with greasy lustre. Madhya Pradesh, India.
8. Blocky and nodular glaebules with a slight surface modification. Sampled from a Latosol derived from eruptiva, Madiun, Java. Some spherical ones are shown.
9. Blocky to fragmental glaebules with smooth surface texture, and spherical ones with smooth and greasy-lustred surface. Sampled from a slag-earth horizon on undulating plateau, Madhya Pradesh, India.
10. Indurated vesicular laterite from Melaka, Malaysia.

- (2) dull brown to greyish yellow, spherical to sub-spherical, with rough surface texture and dull lustre,
- (3) variously colored, blocky, with rough surface texture and dull lustre of slightly modified surface of weathered rock fragments.

The spherical glaeblues with greasy lustre in the first group have been termed pisolite, concretion, buckshot, etc., as mentioned in previous studies. The second group appears bleached with respect to color and surface texture. These two groups show some regular internal structure on splitting, such as concentric bands, the peeling off of the brown crust from the darker core, etc. The third group shows no regular internal structure, and is dominant in samples from Java and the Philippines.

The mangiferous glaeblues are brownish black, spherical and granular, with even to smooth surface texture, and are oblate in shape. The concentric thin laminae are very noticeable on fracture faces. The carbonate glaeblues are irregular and warty in shape, light grey, with rough surface texture and dull lustre; they are sometimes polished and brownish, resembling the bleached ferruginous glaeblues, but easily distinguishable from the latter by effervescence on contact with hydrochloric acid.

Comparison of the chemical composition of the glaeblues with that of the soil matrices indicated the change in the content of each chemical element during glaeblue formation.

Frequency distribution is illustrated for the contents of the nine oxides and for the molar ratio of $\text{TiO}_2/\text{Al}_2\text{O}_3$ (Fig. 3). The soils analyzed comprise four hundred and ten surface soil samples of padi soils in tropical Asia (Kawaguchi and Kyuma, 1974), but they are not necessarily the samples from which glaeblues were separated. They cover, however, a range of chemical composition wide enough to provide a soil matrix for consideration of the loss and gain of elements in glaeblue formation in tropical soils.

The frequency distribution patterns of the $\text{TiO}_2/\text{Al}_2\text{O}_3$ ratio are similar for glaeblues and soils, which may indicate the pedogenic nature of the glaeblue formation.

Relatively high titania content (1.5–2.5%) in glaeblues from Java and the Philippines might be due to the volcanic origin of parent materials, but it is not as high as the titania content reported in ferruginous concretions in Hawaiian Latosol (Sherman and Kanehiro, 1954).

While it is known that active volcanism enhances the $\text{TiO}_2/\text{Al}_2\text{O}_3$ ratio in ocean sediments (Bostrom *et al.*, 1973), this is not the case for surface soil samples. The highest $\text{TiO}_2/\text{Al}_2\text{O}_3$ ratios in the soil samples were for the quartziferous sandy soils of the Khorat plateau of Thailand, but not for soils from active volcanic areas




Fig. 3 Change in chemical composition of glaebules compared to that of soils

■ glaebules ($n=57$). □ soils ($n=410$).

ordinate: frequency as percentage of total number of samples.

abscissa: oxide content in percentage on ignited basis.

of Java and the Philippines, where the high alumina content in soils due to advanced alteration of ferro-magnesian minerals to clay caused rather low $\text{TiO}_2/\text{Al}_2\text{O}_3$ ratios.

The patterns of iron and manganese clearly show high mobility of both elements, which are transported and accumulated as glaebules.

It is interesting to note that calcium and magnesium, which leach rapidly from primary minerals, are found in glaebules to the same extent as in soils. A few peaks appear far away from the mode in case of calcium, while no such peaks appear for magnesium. This difference can be interpreted as indicating that mobilized calcium tends to form free carbonate, whereas magnesium tends to be philic to clay structure.

The very low potassium content of the glaebules is due to the presence of glaebules in the Khorat plateau of Thailand and India. Except for these cases, potassium exists in glaebules, as in soils, at fairly high levels, and this indicates the high resistance of potassium to leaching, even under tropical weathering conditions.

The overall coincidence between glaebules and soil in the frequency distribution patterns of Al_2O_3 , MgO and K_2O , which accumulate in clay minerals, suggests that clay fractions in the vicinity of the precipitation center have been incorporated into glaebules.

The loss of SiO_2 content in glaebules is very outstanding. This is due either to the leaching of silica during intensive weathering to form residual weathering crusts, or to the exclusion of sand grains in plasma precipitation.

Differential precipitation of iron and manganese is frequently observed in micro-scale as exemplified by alternate concentric layers in single glaebules (Table 2). This is also observable in macro-scale, as in the predominance of ferruginous glaebules in highly leached soils on older terraces and that of manganiferous ones in lower-lying base-rich soils.

The differential concentration of either iron or manganese in glaebules as expressed by a ratio (termed the iron-manganese ratio), $\text{Fe}_2\text{O}_3/(\text{Fe}_2\text{O}_3 + \text{Mn}_3\text{O}_4)$ would be positively correlated with the ratio $\text{SiO}_2/\text{Al}_2\text{O}_3$, taking the sum of base content as an internal standard (all contents expressed in percentages). The ratios

Table 2 Chemical composition of outer shell and inner core in manganiferous glaebules
(% on ignited basis)

Sample		SiO_2	Al_2O_3	Fe_2O_3	Mn_3O_4	CaO	MgO	TiO_2	K_2O	P_2O_5	$\text{SiO}_2/\text{Al}_2\text{O}_3$
26-1	outer shell	25.1	8.2	53.0	11.8	0.09	0.85	0.86	0	0.03	5.3
26-2	core	29.8	13.3	22.8	31.6	0.35	0.49	1.59	0	0.02	3.7
41-1	outer shell	33.7	17.6	19.5	23.6	1.43	0.94	1.31	1.19	0.55	3.2
41-2	core	24.6	18.8	17.8	33.0	1.76	1.03	1.35	1.05	0.43	2.2

Notes: 26 is a spherical granular (7–10 mm) manganiferous glaebules with smooth surface texture and dull surface lustre. From the surface horizon of a Latosol on eruptiva, Cauayan, Luzon, the Philippines.

41 is an oblate (7–15 mm) manganiferous glaebule with mammilated surface and dull lustre. From the surface horizon of Reddish-brown soil on coral limestone, Shimajiri, Okinawa, Japan.

are plotted on log-log coordinates in Fig. 4. The positive correlation is thought to be due to the increase of iron relative to manganese content as the silica (presumed to stand for quartziferous sand) increases relatively to alumina (clay). This result accords with the fact that iron oxides are known to be relatively immobile compared

Table 3 Relationship between major chemical composition and content of some minor elements in some ferruginous, manganiferous and carbonate glaebules

(% for oxides and ppm for elements)

Sample	Ferruginous				Manganiferous				Carbonate		
	6-2	15-3	36-1	43	14-1	26-1	42-1	42-2	33-1	34-1	35-1
SiO ₂	30.4	11.1	21.7	20.4	28.9	29.8	28.9	28.6	23.8	30.1	21.8
Al ₂ O ₃	7.9	10.8	10.5	19.5	15.3	13.3	16.9	16.2	5.2	5.1	6.3
Fe ₂ O ₃	60.3	74.7	66.6	58.2	33.7	22.8	22.9	26.0	1.2	1.9	4.0
Mn ₂ O ₃	0.32	0.57	0.62	0.18	18.3	31.6	25.8	23.7	0.41	0.42	0.59
CaO	0	0.12	0.29	0	0.08	0.35	1.45	1.49	66.0	57.8	60.7
MgO	0.32	0.94	0.72	0.97	0.98	0.49	1.16	1.54	2.37	3.52	5.32
TiO ₂	0.47	0.76	0.79	0.64	2.16	1.59	1.39	1.25	0.12	0.20	0.28
K ₂ O	0	0	0.59	0	0	0	1.08	0.99	0.63	0.75	0.73
P ₂ O ₅	0.22	0.90	0.17	0.13	0.47	0.02	0.40	0.48	0.22	0.17	0.20
Na ₂ O	0.057	0.275	0.044	0.020	0.058	0.117	0.102	0.148	0.179	0.222	0.458
Cs	4.9	2.1	2.4	0.9	4.6	4.1	8.0	3.0	2.0	2.8	2.5
Ba	458	283	492	56	1665	5243	2643	1740	390	872	506
Ga	24.5	28.1	(25.0)	55.1	(25.0)	(18.0)	(30.0)	(35.0)	5.8	7.2	(9.0)
Sc	15.5	24.4	18.3	44.2	18.8	14.1	24.5	27.5	4.2	6.5	6.7
La	39.0	11.6	40.5	11.4	44.3	116	73.5	55.7	38.6	162	164
As	171	142	417	845	25.8	13.9	123	131	12.3	21.0	6.0
Sb	10.7	2.6	15.7	9.7	2.28	1.85	11.3	8.05	0.64	0.30	0.53
Cr	382	71.2	246	903	16.9	37.1	85.6	84.7	21.0	28.7	26.5
Co	128	30.4	107	1.2	546	1884	637	723	14.8	23.0	34.2

Note 1. Oxides are on ignited basis, and elements are on air-dry basis. Values for Ga are estimated using the proportional relationship between Ga and Sc. Values in parenthesis for Ba are relative values.

Note 2. Description of samples.

6-2. Spherical granular (2–5 mm) ferruginous glaebules with smooth surface texture and dull lustre. Sampled from subsoil of Low Humic Gley soil on old alluvium in old delta, Thailand.

15-3. Spherical granular (2 mm) ferruginous glaebules with rough surface texture and dull lustre. Sampled from sub-surface horizon of Alluvial soil, Tangerang, Java.

36-1. Spherical granular (3 mm) ferruginous glaebules with smooth surface texture and greasy lustre. Sampled from surface horizon of local alluvium in Red and Yellow soil zone on undulating plateau, Madhya Pradesh, India.

43. Indurated vesicular laterite derived from saprolitic argillaceous rocks in Melaka, Malaysia.

14-1. Spherical granular (5–8 mm) manganiferous glaebules with smooth surface texture and dull lustre. Sampled from subsoil of a Latosol derived from basic and intermediate eruptiva, Bogor, Java.

26-1. See note Table 2.

42-1. Inner core of manganiferous glaebules with mammillated surface texture from Okinawa, Japan. Similar to 41. See note Table 2.

42-2. Outer shell of the above glaebule.

33-1. Subspherical, warty carbonate glaebule (1.5–8 mm) with rough surface texture. Sampled from subsoil of a Gray and Brown soil impregnated with salts on a low terrace in the Ganges South Delta, Uttar Pradesh, India.

34-1. Subspherical, warty carbonate glaebules (1.5–15 mm) with rough surface texture. Sampled from the sub-surface horizon of a Gray and Brown soil impregnated with salts on a high terrace in the Ganges South Delta, Uttar Pradesh, India.

35-1. Spherical (8–10 mm) carbonate glaebules with rough surface texture. Sampled from surface horizon of Red Loam on local alluvium on an undulating plateau, Orissa, India.

to manganese oxides.

The correlation was even greater when the abscissa was replaced by a ratio $\text{SiO}_2/(\text{CaO} + \text{MgO} + \text{K}_2\text{O})$. This might be interpreted as indicating that iron may be concentrated in combination with colloidal silica, which may act as a protective colloid for iron, since in an acidic soil pH range, silica and poorly crystallized manganese oxides are known to be negatively charged, and iron oxides positively charged.

The content of some minor elements, determined by neutron activation analysis for a few glauabules (Table 3), seems to indicate that each element is selectively accumulated mainly depending on its charge relative to that of either iron and manganese oxides sol (Landergrén, 1948). Chromium and arsenic clearly accumulate in ferruginous glauabules, while cobalt, barium and lanthanoids accumulate in mangiferous ones. The high correlation between gallium and aluminium content can be explained by Goldschmidt's hypothesis concerning ionic radii (1945).

The accumulation of phosphorus in glauabules is noticeable (Fig. 4). However, no significant correlation between phosphorus and one or a combination of iron, aluminium, or manganese content is found. Phosphorus is a constituent of such primary minerals as apatite which is abundant in volcanic rocks, and this may be the




Fig. 4 Differential accumulation of iron and manganese in relation to the Silica/Alumina ratio. Base content of the glauabules was used as an internal standard.
ordinate: $\text{Fe}_2\text{O}_3(\%)/(\text{Fe}_2\text{O}_3 + \text{Mn}_2\text{O}_3)(\%)/(\text{CaO} + \text{MgO} + \text{K}_2\text{O})(\%)$
abscissa: $\text{SiO}_2(\%)/\text{Al}_2\text{O}_3(\%)/(\text{CaO} + \text{MgO} + \text{K}_2\text{O})(\%)$

cause of the high phosphorus content of glaebules and soils from Java.

2. Mineralogical composition

Some representative diffractograms of glaebules are shown in Figure 5.

Goethite was the predominant iron mineral in most ferruginous glaebules. Haematite was dominant only in the glaebules with laterite fabric and in the indurated laterite. Thus, glaebules with laterite fabric were considered to have been derived from the disintegrated laterite crust.

Calcite was the predominant mineral in carbonate glaebules.

Although goethite was detected, with rather diffuse peaks, no manganese minerals were detectable in manganiferous glaebules.

Quartz was almost completely absent in glaebules characterized by a low $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio and by a relative abundance of haematite, while it was common to abundant in pisolitic concretions with a high $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio.

Feldspars were found in the samples from the northern intermountain basins of Thailand, and also in the immature glaebules found in volcanic regions of the Philippines and Java.

Gibbsite was not detectable even in the samples with a low $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio. The clay minerals have not been broken down intensively that the alumina released




Fig. 5 Crystalline minerals in glaebules. Diffractograms from powdered specimen with Fe K_α . G: goethite, H: haematite, F: feldspar, Q: quartz, C: calcite. From top to bottom: carbonate (33-1, see Table 3), ferruginous (4, see Table 4), manganiferous (41-1, see Table 2), ferruginous (2, see Table 4), ferruginous (38-3, see Table 4), indurated laterite (43, see Table 3).

is accumulated in the form of gibbsite. This agrees well with the general trend, to be stated later, that the clay minerals in glaeboles and in soils are of nearly the same composition.

The color of the ground powder can be related to the nature of the iron minerals and manganese content. Goethite rich ferruginous glaeboles are bright brown to dark brown (7.5YR to 10YR in hue and value/chroma ranging from 5/6 to 3/4). As manganese increases the value and chroma decrease to 3/3–2/3, and if there is a high haematite content the hue shifts to 5YR or redder. Manganiferous glaeboles mostly show brownish black (7.5YR 2/2). Haematite-rich iron glaeboles are dark reddish brown to dark red in color (10R3/3–3/6).

It was previously claimed by Harada (1937) that an oxalic acid-potassium oxalate solution of pH 4.1 was effective in differential dissolution of iron compounds. Therefore, his method was adopted to study the present samples, with slight modifications, as follows: 100 mg of ground powder was dissolved in 250 ml of 0.1 M potassium oxalate-0.025 M oxalic acid under an UV lamp or in the dark. The extracted iron expressed as Fe_2O_3 was called FeI and FeIII for the light and dark treatment, respectively. A similar dissolution under an UV lamp and using a more dilute solution, 0.015 M potassium oxalate-0.05 M oxalic acid, was carried out to determine FeII. The amount of iron extracted was determined by atomic absorption spectroscopy.

Although the differences (FeI–FeII) and (FeII–FeIII) were assumed by Harada to correspond to the amounts of haematite and limonite, respectively, this was not verified by the results of X-ray diffraction. This was because well crystallized haematite and goethite were only partly dissolved in the solution, even under an UV lamp. The same phenomenon was observed in Mehra-Jackson's deferration treatment.

The ratio FeIII/FeI seems to be better correlated with the degree of crystallization of iron compounds; when the ratio is high, X-ray amorphous iron compounds predominate, and when it is low, crystalline compounds are dominant (Table 4).

The FeIII/FeI ratio seems to be equivalent to the "Aktivitätsgrad" (Schwertmann, 1964) of iron oxides, which was effectively used in combination with the "crystallinity ratio" by Nagatsuka (1972) to distinguish some relict soils (paleosols) from the modern soils in Japan. In this case, however, a decrease in iron solubility for highly crystalline minerals would give a rather low "crystallinity ratio".

Diffraction patterns of clay separates from some glaeboles are shown in Fig. 6. It is interesting to note that the goethite peak persists after Mehra-Jackson's deferration treatment in well crystallized samples.

Clay minerals with 14 to 15 Å spacing on magnesium saturation are vermiculite

Table 4 Relation between iron minerals present and iron compounds extracted by oxalate solution

Sample	Relative abundance		FeIII/FeI %	(FeI – FeIII)/FeT %
	Goe.	Hae.		
38-3	++	++	2.4	71.6
43	+	++	2.8	64.4
2	++	+	6.0	57.3
14-2	++	±	14.3	64.7
15-3	++	±	8.4	61.3
36-2	++	±	3.5	73.1
3	++	—	13.7	58.1
6-2	++	—	3.7	77.1
7-2	++	—	5.4	73.9
26-3	++	—	12.8	68.0
4	+	—	12.1	80.2
7-1	+	—	13.2	77.7
36-3	±	—	36.7	45.2
41-2	±	—	54.1	31.8
5	—	—	38.4	55.2
41-1	—	—	85.2	10.1

Note 1. FeIII/FeI and (FeI–FeIII)/FeT would be equivalent to the “Aktivitätsgrad” of Schwertmann (1964), and to the “crystallinity ratio” of Nagatsuka (1972) respectively.

Note 2. Description of samples.

38-3. Spherical (1.5–5 mm) ferruginous glaebules with smooth surface texture and greasy lustre. Sampled from slag-earth horizon in Red and Yellow soil zone, Madhya Pradesh, India.

43. See note Table 3.

2. Botryoidal spherical ferruginous with greasy lustre coalesced by cementing material. Sampled from indurated hard pan underlying sandy soils on old alluvium, Ubon, Thailand.

14-2. Spherical (2 mm) ferruginous glaebules. See note Table 3.

15-3. See note Table 3.

36-2. Spherical (4–7 mm) ferruginous glaebules. See note Table 3.

3. Spherical (2–6 mm) ferruginous glaebules with smooth surface texture and dull lustre. Occur as ironstone layer in Low Humic Gley soil on old alluvium in intermontane basin, Chiangmai, Thailand.

6-2. See note Table 3.

7-2. Spherical (1.5–4 mm) ferruginous glaebules with smooth surface and greasy lustre. Sampled from throughout a profile of Low Humic Gley soil on old alluvium in old delta, Manom, Thailand.

26-3. Spherical (2–6) with smooth surface and dull lustre. See note Table 3.

4. Spherical (2–4 mm) ferruginous glaebules with rough surface and dull lustre. Sampled from surface to sub-surface horizon of Low Humic Gley soil on semi-recent alluvium in intermontane basin, Chiangmai, Thailand.

7-1. Ferruginous and manganiferous glaebules coalesced into botryoidal ones (5–8 mm). See note for 7-2.

36-3. Inner core of manganiferous glaebules. See note for 36-2.

41-2. See note Table 2.

5. Warty ferruginous glaebules with bleached appearance. Sampled from subsoil of the same profile as 4.

41-1. See note Table 2.

and montmorillonite. 10 Å spacing is due to mica and illite. 7 Å spacing is mostly due to kaolin minerals as this shifts to 10 to 10.5 Å on hydrazin intercalation (Wada and Yamada, 1968).

It should be noted that the clay mineral composition of the clay separates of glaebules is not very different from that of surrounding soils.

For example, in the glaebule samples from Northern Thai intermontane basins




Fig. 6 Crystalline clay minerals in glaebules. Diffractograms from oriented specimen with Cu K α

K-Hyd.: potassium-saturated and intercalated by hydrazin,

K-Ad.: potassium-saturated and air-dried,

Mg-Gly.: magnesium-saturated and glycerated,

Mg-Ad.: magnesium-saturated and air-dried.

1. Kaolin, mica and quartz are dominant. Ferruginous glaebules 4. see Table 4.
2. Vermiculitic montmorillonite interlayered, quartz and kaolin are dominant. Carbonate glaebules 34-1. see Table 3.
3. Kaolinite predominates. Goethite peak is visible at 4. 18 Å. Indurated vesicular laterite. 43. see Table 3.
4. Kaolinite predominates and mica is present. Goethite peak is visible. Ferruginous glaebules 38-1. see Table 4.

(Fig. 6-1), 10 Å spacing due to mica and illite is greatly enhanced relative to that of the soil, whereas the potassium content of the glaebules is significantly less than that in the bulk soil. This can be interpreted as meaning that the glaebules had been formed in the course of weathering of granitic rocks prevalent in the area before they were transported to the detritus slope and from there to the alluvial plain. This may also hold true for pisolitic concretions from the upper Central Plain of Thailand.

In samples from Java and the Philippines no significant difference between glaebules and the soils was detected.

The interesting clay mineral composition of one carbonate glaebule (Fig. 6-2) is to be noted. The sharp and strong peaks at 14 Å to 15 Å of Mg-saturated clay showed a decline to 10 Å spacing on K-saturation, and on glyceration expanded to around 20 Å, with part remaining at 14 Å. Thus the presence of vermiculite and montmorillonite is evident, but the montmorillonite seems to be a grade intermediate to vermiculite and could be called vermiculitic montmorillonite, and interstratified with Al-interlayered mineral (Hattori, private communication). According to Hattori, this kind of montmorillonite is frequently observed in montmorillonitic soils in tropical Asia, except in grumusols where it behaves more ordinarily.

In the indurated laterite crust (Fig. 6-3), kaolinite is the dominant clay mineral. The clay material filling the vermicular voids has almost the same composition as clay material in the crust, except that goethite is lacking.

In glaebules with laterite fabric haematite and goethite are abundant, and kaolinite is much more prominent than clay in the soil.

In conclusion, it appears that the discrepancy in mineral composition of clay fraction between glaebules and soil matrix may be greater or lesser according to the lag between the time of formation of glaebules and soil. When a glaebule is an *in situ* formation, the discrepancy may be small, as could be the case for most manganese glaebules from Central Luzon and East-Central Java. When it is transported, the discrepancy may be greater; younger minerals may be found in glaebules as is the case with ferruginous glaebules from Northern and Central Thailand, and minerals at advanced stages of weathering may be found as is the case with laterite-derived glaebules from Madhya Pradesh, India.

3. Micromorphological observations

The most striking feature of the thin-sections of glaebules was observed under reflected light; the fabric and plasma were clearly differentiated.

The fabric of indurated laterite from Malaysia is characterized by a mosaic of the following two areas; (1) one is that part apparently corresponding to the vesicular void space, which shows a radial plasma orientation appearing brownish under the

transmitted light and bright yellow (abbreviated hereafter as brown/bright yellow), and the other is the area of the hardened crust, in which, while showing a similar orientation, plasma tends to be granulated into relatively coarse granules appearing opaque/lustrous red. Granules are “tied” (Humbert, 1948), or “knitted” (Alexander and Cady, 1962) by channels filled with clay. The iron compound is considered to be amorphous in the vesicular void, and haematite and goethite in the granulated area. The fabric described above may correspond to the “Braunlehm-Laterite” of Kubiena (1970).

The similarity of the two areas in both clay mineral composition and in orientation pattern may be taken as evidence of *in situ* plasma segregation. The growth of the network of granules contributed to the hardening of the laterite. Another characteristic of laterite fabric is the absence of coarse quartz grains.

The Braunlehm-Laterite fabric is also found in several glaebules from lowland soils of India and of Northeast Thailand, regardless of their shape, which may be concretionary, botryoidal, or nodular. The thin sections show that the laterite fragment is coated and/or coalesced by thinly laminated plasma, resulting in different shapes. There is little doubt that disintegrated laterite fragments transported to low-lying areas acted as cores which absorbed plasma from the surrounding soil matrix under the alternate dry and wet conditions. The fact that fragmental slag-earths on the higher part of undulating terrain in Madhya Pradesh, for example, change into spherical glaebules in local alluvia suggests that the spherical shape has resulted from abrasion during transportation.

In the laterite fabric the size of the individual lustrous red granules ranged

Fig. 7 Micromorphology of some representative glaebules. Scale 0.5 mm

1. Indurated vesicular laterite. $\times 28$. Reflected light. Vesicular voids are filled by yellow clay (light-toned part). Dark granules are haematite crystallites. White part is vacant void.
2. Same sample at higher magnification. $\times 70$. Transmitted light (parallel nicols).
3. Ferruginous glaebule with earthy braunlehm fabric. $\times 28$. Reflected light. Core of earthy braunlehm fabric is coated with concentric thin laminae of goethite and clay.
4. The same field under crossed nicols. Bright spots are quartz grains.
5. Receptized ferruginous glaebule. $\times 70$. Reflected light. The inner part of the core remains lateritized earthy braunlehm fabric, while the outer part is receptized to braunlehm. The core is coated with laminae of goethite and clay. Spots are quartz grains.
6. Concentric manganiferous glaebule. $\times 28$. Reflected light. Quartz grains are cemented by iron oxides, on which alternate precipitation of manganese (darker laminae) and iron (lighter laminae) oxides is apparent. Outermost coating is ferruginous.
7. Poorly differentiated manganiferous glaebule. $\times 28$. Reflected and transmitted light (parallel nicols). Quartz nest in the center with unlaminated manganiferous precipitated on it.
8. Poorly differentiated glaebule with weathered rock fabric. $\times 28$. Reflected and transmitted light (parallel nicols). At top and bottom argillized feldspar can be seen; augite (pointed grains) and hornblende (columnar grains) are embedded in braunlehm.
9. Ferruginous glaebule with weathered rock fabric. $\times 28$. Transmitted light (crossed nicols). Striated extinction pattern is derived from weathered volcanic rock.
10. Carbonate glaebule. $\times 28$. Reflected and transmitted light (parallel nicols). Microcrystals of calcite densely fill the groundmass.




Fig. 7

from 10 to 30 μ (Fig. 7-2). These granules appear to be identical with the “droplet” reported by Hamilton (1964), which was, however, exceptional in the old iron crust of west Africa. Our contention is that the segregation of iron compounds into granules or droplets of haematite is a rather common phenomenon in the laterite formations of tropical Asia, and that the glaebules derived from laterite can be easily distinguished from other kinds of glaebules. The second type of fabric, one of the commonest, is found in the pisolitic glaebules from the upper Central Plain of Thailand and from India (Fig. 7-3, 4). They are characterized by a fairly thick coating of thinly laminated plasma layers around the inner core. The coating is reddish brown to opaque/yellow to dark yellowish brown, almost non-birefringent, has few voids other than some cracking voids, and has few quartz grains; the inner core is variously colored, with varying amounts of quartz grains but definitely more coarse grains than in the coating, and also has many voids mostly lined by birefringent plasma.

The fabric of the inner core may be understood as a variety of stabilized, that is, “erdified”, braunlehm in which the dense groundmass of braunlehm is transformed into a structured groundmass, and the iron hydroxides move toward the crystallization center along or in the voids. The degree of crystallization of the iron hydroxides may vary.

Based on the degree of “encasing” (Kubiena, 1970) of the groundmass by threads of plasma, and also on the degree of crystallization of plasma precipitated on ped faces or in the groundmass, several intergrades would occur, ranging from earthy braunlehm, via *erde*, to lateritized earthy braunlehm.

As described above, a very marked contrast exists between the coating and the inner core; absolute accumulation of iron from the transporting solution no doubt occurs in the former, while a relative accumulation of sesquioxides would be the main process occurring in the precursor of the core.

Since the fabric of the soil matrix in a low humic gley soil of, for example, the upper central plain of Thailand is dominated by earthy braunlehm fabric, those glaebules with lateritized earthy braunlehm fabric in the inner core have most probably developed on ferruginous soil clods transported from higher terrain. *In situ* formation is conceived, however, for those which have many quartz grains cemented by manganese compounds in the inner core and the laminated ferruginous shells in the coating.

The red mottles found in lowland alluvial soils may have been transformed into harder concretions or nodules as the sediments dried out, either because of uplift of the terrain or a lowering of the water table associated with an eustatic change. Hardening of red mottles is observed even today when subsoils are dug out and

exposed to the surface conditions. Then they give nodular hardened materials with vesicular voids filled with clay, very closely resembling laterite fragments. But materials thus hardened can be distinguished from transported glaebules by such features as the lack of smooth surface texture and spherical shape, and the absence of a laminated coating of plasma.

The third type of fabric can be seen in the glaebules with a bleached appearance (Fig. 2), which have a fabric suggestive of a change of the environment (Fig. 7-5). Repetition of iron compounds causes differentiation of the formerly homogeneous fabric into two parts; the inner part presumably consists of the former fabric of laterite and lateritized earthy braunlehm, whereas the outer part has changed into braunlehm. The components of the fabric except for the type of plasma, for example, the distribution of skeletal grains and voids, remain unchanged in parts.

This may be interpreted as follows; the ferruginous glaebules which had been formed in higher terrain with favorable conditions for dehydration and crystallization of precipitated and/or segregated iron compounds were transported to low-lying areas where redissolution or rehydration of the iron compounds took place under reduced conditions that were prevalent during the submerged period. The dissolution of hard iron concretions has also been reported by Nye (1954) as occurring in soil on the down-slope and valley bottom of a West African catena.

Redissolution of precipitated plasma would be a more common phenomenon in manganiferous glaebules since the many forms of manganese compounds are of higher solubility and more mobile than iron compounds. In fact, the manganiferous glaebules in weakly acid to acid soils are in many cases coated by an often very thin ferruginous crust; otherwise, the development of the glaebules would have been halted by the redissolution process. Thus the fabric of manganiferous glaebules is characterized by alternate precipitation of manganese and iron compounds, resulting in thinly laminated layers of remarkably developed concentric pattern (Fig. 7-6).

In calcareous soils areas, for example, the Ganges Plain of India where manganiferous and ferruginous glaebules coexist with carbonate glaebules, this concentricity is not prominent in manganiferous glaebules (Fig. 7-7). This implies a decline in the dissolution of iron and manganese compounds in an alkaline medium, while their precipitation from the transporting solution is favored.

In view of amorphous nature and high solubility of the precipitated iron compounds in the concentric fabric, many manganiferous glaebules seem to be of *in situ* formation.

The fifth type of fabric is seen in many glaebules from Java and the Philippines,

containing high amounts of weatherable minerals, such as plagioclase, augite, hornblende, titaniferous minerals, and sometimes glass, all of volcanic origin. The degree of crystallization of precipitated compounds is low, as revealed by X-ray diffraction analyses. The fabric is predominantly of earthy braunlehm type, and plasma separation is mostly confined to micro-scale segregation from weatherable minerals embedded in the groundmass. (Fig. 7-8)

In other words, these glaeboles show an initial stage of redistribution of plasma within a rock fragment rich in iron, manganese and bases. Part of the diffused plasma is precipitated on the surface of weathered rock fragments, and sometimes make them varnished (Fig. 2-8). Other parts of the diffused plasma would be carried farther by the transporting solution finally to precipitate on incipient glaeboles. The varnished rock fragment, too, would be transported physically to low-lying land, and become the precursor of a glaebole. The spherical ferruginous glaeboles with a relatively thick coating found in the older alluvium of Cagayan valley, Luzon (Fig. 7-9) are probably formed in this way. The inner core has a relict of rock fabric. The striated extinction pattern seen under crossed nicols is associated with the argillized rock structure of volcanic origin.

The fabric of the carbonate glaeboles is characterized by abundant microcrystals, densely scattered through the whole groundmass (Fig. 7-10). These microcrystals are very highly birefringent and from powder diffraction data are considered to be calcite. It should be noted that calcite microcrystals are sometimes found in other types of glaeboles at the same site, and these microcrystals show a referred distribution related to voids and the concentric layering pattern.

4. Conclusion

Following the results and discussions presented in the preceeding section, it may be possible and of interest to classify glaeboles by their properties. We prefer a classification that allows us, first of all, to separate these glaeboles of *in situ* formation from those formed otherwise. A rigorous distinction, however, between autochthonous and allochthonous formation is difficult because of the transportation of materials both chemically and physically. Autochthonous formation, therefore, is defined as the mobilization and precipitation of plasma that is mostly from the same geomorphologic terrain to form glaeboles.

The following are considered autochthonous formations.

- (1) Oblate to spherical manganiferous glaeboles, Those with smooth surface mostly occur in low-lying base-rich soils, particularly in the montmorillonitic soils of Java and the Philippines. Those with slightly mammilated surface texture tend to occur in the base-rich soils on intermediate to basic eruptiva and on lime-

stone. Alternate precipitations of amorphous manganese and iron oxides in a concentric pattern is a prominent feature.

- (2) Spherical ferruginous glaebules with a thick manganiferous core, which occur mainly on natural levees and on low terraces.
- (3) Warty and irregular carbonate glaebules which mainly occur in limestone areas, carbonate-impregnated lowlands, and on volcanic fans of calc-alkali rocks. Microcrystals of calcite densely fill groundmass.
- (4) Blocky to irregular weathered rock fragments with a slight modification of surface texture. Micro-scale segregation of iron and manganese occurs in association with partly weathered mineral grains such as pyroxenes, amphiboles and titaniferous minerals. These glaebules mostly occur on volcanic fans in Java and the Philippines.

The following may be grouped as autochthonous transformations on allochthonous glaebules.

- (5) Subspherical to spherical ferruginous glaebules with light greyish color and dull lustre. The outer part consists of repetized iron compounds, while the inner part is a laterite or lateritized fabric. Sometimes they are coated by concentric laminae. These occur on fan-terrace complex areas, and on local alluvia on older terrains.
- (6) Spherical, botryoidal and nodular ferruginous glaebules with greasy surface lustre and smooth texture. The core with laterite or lateritized fabric is coated with concentric ferruginous laminae. They occur on fan-terrace complex areas and on local alluvia on older terrain.
- (7) Glaebules with a similar appearance to the above, but with a rock fabric in the core, occurring mostly on the older alluvia of the Luzon Plain and Java.

All the varieties of glaebules are listed in Table 5, in which further subdivision of the above-mentioned glaebules is included, putting a tentative name to each.

Iron, manganese, and calcium are particularly concentrated in glaebules compared to the content in soils. This shows clearly that they can be mobilized, transported and precipitated. The decisive processes involved in the geographical redistribution of these elements are the chemical transformation of these elements and the hydrological behavior of the transporting interflow (Milot, 1970).

In addition, the physical transport of glaebules, which is clear from the present study, should be stressed. Physical transport may be carried out by soil creep, by sheet erosion, or by dissection of older terrain, in which glaebules are removed from higher land and transported to lower land. An intricate balance between these two processes leads to differential distribution of specific glaebules on a specific terrain.

Table 5 Classification of glaeboles found in some tropical lowland soils

Varieties	Appearance	Chemical composition			Iron-Manganese ratio	SiO ₂ /Al ₂ O ₃
		Fe ₂ O ₃ (%)	Mn ₃ O ₄ (%)	CaO (%)		
I Autochthonous formations						
1. Concentric manganiferous glaeboles.	a. spherical, smooth surface.	18-34	15-33	0.08-3.3	0.3-0.6	3-7
	b. oblate, mammillated surface.					2-3
2. Poorly differentiated manganiferous gl.	oblate, smooth surface.	15-53	3-10	0.3-0.6	0.7-0.8	5-10
3. Ferruginous gl. with manganiferous core.	spherical, smooth surface with greasy lustre.					
4. Carbonate gl.	warty, blocky to subspherical.	1.2-26	0.4-14	13-66	0.66-0.82	5-10
5. Poorly differentiated gl. with weathered rock fabric.	blocky, rough surface, dull lustre.					
II Autochthonous transformation of allochthonous glaeboles						
6. Repeptized ferruginous gl.	spherical to subspherical, smooth to dull surface, dull lustre.	27-68	3≥	0.7≥	0.90-0.997	1.7-2.9
7. Ferruginous gl. with laterite fabric.	spherical, nodular to botryoidal, smooth surface with greasy lustre.					higher
8. Ferruginous gl. with lateritized fabric.	spherical, smooth surface with dull to greasy lustre.					1.7-2.1
9. Ferruginous gl. with rock fabric.	spherical to subspherical, smooth surface with dull to greasy lustre.					
Mineralogical features		Micromorphological features		Occurrence		
1a. Go. (-) - (±), Qu. (+) - (++)	manganese minerals.	alternate manganiferous and ferruginous laminae developed concentrically throughout.		base-rich soils on low terrace, recent alluvium. base-rich soils on eruptiva and limestone.		
2.						
3. Go. (++)	Qu. (++)	concentric manganiferous core, with concentric iron coating.		soils on natural levee and low terrace.		
4. Calc. (++)	Qu. (±) - (++)	micro-crystals of calcite densely filling the matrix.		carbonate-impregnated areas, volcanic fan of calc-alkali eruptiva.		
5. Fl. (+) - (++)	negative on Go.	micro-segregation of the plasma associated with heavy minerals (Py, Am, titaniferous minerals).		soils on volcanic fan, Java and Luzon.		
6. Ha. (+) - (++)	Go. (++)	iron oxides repeptized on outer part, laterite and lateritized fabric within.		Low Humic Gley soils on fan-terrace complexes.		
7. Ha. (++)	Go. (++)	iron oxides granulated into coarser crystallites, lehm filling the channels.		local alluvium and residual sites on undulating plateau.		
8. Ha. (+)	Go. (++)	'threads' of iron oxides encasing matrix, concentric laminae coating the core.		Low Humic Gley soils on fan-terrace complex (middle terrace).		
9. Go. (++)		weathered rock fabric in the core, concentric laminae coating.		soils on old alluvium in volcanic areas.		
Key;						
gl.	glaebole	(-)	absent			
Go.	goethite	(±)	just detectable			
Ha.	haematite	(+)	present			
Qu.	quartz	(++)	common			
Fl.	felspar	(+++)	profuse			
Calc.	calcite					

Summary

The chemical and mineral composition, and micromorphological properties of several kinds of glaebules collected from padi soils in South and Southeast Asian countries were studied.

They were separated into ferruginous, manganiferous and carbonate glaebules, in terms of chemical composition. Silica content decreased in glaebules in comparison with that of the padi soils of the area, while the content of bases remained at almost the same level in glaebules and soils. Alumina, titania, and the molar ratio $\text{TiO}_2/\text{Al}_2\text{O}_3$ also stayed at the same level in both. These results suggest that clay in the vicinity of the precipitation center was incorporated into glaebules, while sand grains were excluded.

The relative concentration of iron to manganese was correlated with the $\text{SiO}_2/\text{Al}_2\text{O}_3$ ratio, which could be an index of the degree of weathering. This result is compatible with observations that ferruginous glaebules are frequently observed on highly leached soils on older terraces.

The most common iron mineral was goethite except in glaebules with a laterite fabric, in which haematite predominates.

The crystalline structure of iron minerals estimated by X-ray diffraction appeared to be well correlated with a ratio nearly equivalent to the "Aktivitätsgrad". Well crystallized goethite and haematite resisted Mehra-Jackson's deferration treatment.

The clay mineral composition of many glaebules was more or less similar to that of the matrix soil, but the discrepancy was considerable for glaebules thought to have been transported.

Laterite fabric is characterized by the growth of iron oxides into coarse granules, such granulated areas being connected or demarcated by channels filled with bi-refrangent clay. Several ferruginous glaebules with laterite fabric were found among the samples from India and Northeast Thailand. They were embedded in the soils by physical transportation of laterite fragments, and underwent subsequent transformation of fabric as exemplified by a thin concentric coating, or by repetization of the outer part. The manganiferous glaebules have clearly developed concentric laminae, indicating the alternate precipitation of manganese and iron oxides *in situ*.

The various glaebules were classified into two groups by their properties, with the aim of distinguishing between autochthonous and allochthonous formations.

References

- Alexander, L. T. and J. G. Cady. 1962. "Genesis and hardening of laterite in soils," *Soil Cons. Serv. Tech. Bull.* 1282, pp. 90.
- Barrera, A. 1964. "Classification and utilization of some Philippine soils," *J. Trop. Geography*, 18, pp. 17-29.
- Boström, K., T. Kraemer and S. Gartner. 1973. "Provenance and accumulation rates of opaline silica, Al, Ti, Fe, Mn, Cu, Ni, and Co in Pacific pelagic sediments," *Chem. Geol.*, 11, pp. 123-148.
- Brewer, R. 1964. *Fabric and Mineral Analysis of Soils* (Pedological features III, Glaebules), John Wiley & Sons, Inc.
- Drosdoff, M. and C. C. Nikiforoff. 1940. "Iron-manganese concretions in Dayton soils," *Soil Sci.*, 49, pp. 333-345.
- Fridland, V. M. 1964. *Soils and Weathering Crusts of Humid Tropics*, Nauka, Moscow. (in Russian)
- Goldschmidt, V. M. 1945. "The geochemical background of minor-element distribution," *Soil Sci.*, 60, pp. 1-7.
- Hamilton, R. 1964. "A short note on droplet-formation in ironcrusts," ed. by A. Yongierius, *Soil Micromorphology*. Elsevier Publ. Co.
- Harada, H. 1937. "Studies on the weathering of igneous rocks, Part IV, Determinative method of free iron oxides (haematite and limonite) by photochemical reduction," *Nihon Nokei Kagaku Kai Shi*, 13, pp. 383-391.
- Humbert, R. P. 1948. "The genesis of laterite," *Soil Sci.*, 65, pp. 281-290.
- Kawaguchi, K. and K. Kyuma. 1974. "Paddy Soils in Tropical Asia, Part 2—Description of fertility characteristics," *Tonan Ajia Kenkyu*, Vol. 12, No. 2, pp. 177-192.
- Kellogg, C. E. 1949. "Preliminary suggestion for the classification and nomenclature for Great Soil Groups in tropical and equatorial regions," *Comm. Bur. Soil Sci. Tech. Commun.*, 46, pp. 76-85.
- Kubiena, W. L. 1970. *Micromorphological Features of Soil Geography*. Rutgers University Press.
- Landergrén, S. 1948. "On the geochemistry of Swedish iron ores and associated rocks," *Sveriges Geol. Under.*, Ser. C, 42(5), 112-124, 129-138. Reprinted in 1975. *Geochemistry of Iron*, ed. by H. Lepp, Dowden: Hutchinson & Ross, Inc.
- Matsuzaka, Y., H. Otowa, Y. Yamada and T. Hamazaki. 1971. "Soils of Okinawa and Kume Islands," *Bull. National Inst. Agr. Sci. (Japan)*, Series B, 22, pp. 305-404.
- Mehra, O. P. and M. L. Jackson. 1960. "Iron oxide removal from soil clays by a dithionite-citrate system buffered with sodium bicarbonate," *Clays and Clay Minerals*, New York: Pergamon Press, pp. 317-327.
- Millot, G. 1970. "Weathering and soil clays," *Geology of Clays, Weathering, Sedimentology, Geochemistry*, translated by W. R. Farrand and H. Paquet, Springer Verlag.
- Moormann, F. R. and S. Rojanasoonthon. 1967. *General Soil Conditions of Thailand*, Scale 1:2,500,000.
- Nagatsuka, S. 1972. "Studies on genesis and classification of soils in warm-temperate regions of south-west Japan, Part 3, Some features in distribution and mode of existence of free iron and aluminium oxides in the soil profile," *Soil Sci. Plant Nutr.*, 18, pp. 147-154.
- Norrish, K. and B. W. Chappell. 1967. "X-ray fluorescence spectrography," *Physical Methods in Determinative Mineralogy*, ed. by J. Zussman, Academic Press.
- Nye, P. H. 1954. "Some soil-forming processes in the humid tropics, I. A field study of a catena in the West African forest," *J. Soil Sci.*, 5, pp. 7-21.
- Okagawa, N. 1970. "Geomorphological analyses of the central plain of Luzon in relation to the characteristics of soil sampling sites," *Tonan Ajia Kenkyu*, Vol. 8, No. 1, pp. 113-122.
- Schwertmann, U. 1964. "Différenciation der Eisenoxide des Bodens durch Extraktion mit Ammonium-oxalat-Lösung," *Z. Pflanzenernähr. Dung. Boden.*, 105, pp. 194-202.
- Sherman, G. C. and Y. Kanehiro. 1954. "Origin and development of ferruginous concretion in Hawaiian latosols," *Soil Sci.*, 77, pp. 1-8.
- Suprpto-hardjo, M., D. Z. Sahertian, and R. Dudal. 1960. *Exploratory Soil Map of Java and Madura*, Scale 1:1,000,000, Bogor: Soil Research Institute.
- Survey of India Offices, 1954. *Soil Map of India*, Scale 1:4,435,200.
- Takaya, Y. 1968. "Quaternary outcrops in the central plain of Thailand," ed. by K. Takimoto, *Geology and Mineral Resources in Thailand and Malaysia*, Kyoto: The Center for Southeast Asian Studies.
- Takaya, Y. and K. Kyuma. 1968. "Geomorphological setting of the soil sampling sites in India," *Tonan Ajia Kenkyu* Vol. 6, No. 3, pp. 614-625.
- Van Wambeke, A. R. 1962. "Criteria for classifying tropical soils by age," *J. Soil Sci.*, 13, pp. 124-132.
- Wada, K. and H. Yamada. 1968. "Hydrazin intercalation-intersalation for differentiation of Kaolin minerals from chlorites," *Am. Miner.*, 53, pp. 334-339.